

Nonlinear Transformation Group of CAR Fermion Algebra

Mitsuo Abe^{a)} and Katsunori Kawamura^{b)}

Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan

Abstract

Based on our previous work on the recursive fermion system in the Cuntz algebra, it is shown that a nonlinear transformation group of the CAR fermion algebra is induced from a $U(2^p)$ action on the Cuntz algebra $\mathcal{O}_{2^p}$ with an arbitrary positive integer p . In general, these nonlinear transformations are expressed in terms of finite polynomials in generators. Some Bogoliubov transformations are involved as special cases.

^{a)}E-mail: abe@kurims.kyoto-u.ac.jp

^{b)}E-mail: kawamura@kurims.kyoto-u.ac.jp

In our previous paper,¹⁾ we have introduced the recursive fermion system (RFS_p) which gives embeddings of CAR into the Cuntz algebra²⁾ $\mathcal{O}_{2^p}$ with an arbitrary positive integer p . As for the special case, which we call the *standard* RFS_p, CAR is embedded onto the $U(1)$ -invariant $*$ -subalgebra $\mathcal{O}_{2^p}^{U(1)}$ of $\mathcal{O}_{2^p}$. Here, the $U(1)$ action on $\mathcal{O}_d$ is defined by

$$\gamma_z : s_i \mapsto z s_i, \quad z \in \mathbf{C}, \quad |z| = 1 \quad (1)$$

with s_i ($i = 1, \dots, d$) being the generators of $\mathcal{O}_d$. Since an automorphism of $\mathcal{O}_{2^p}$, which is described by a $U(2^p)$ action, also keeps $\mathcal{O}_{2^p}^{U(1)}$ invariant, it induces a transformation of CAR. We find this type of transformations is *nonlinear* with respect to generators $\{a_m, a_n^* \mid m, n = 1, 2, \dots\}$ in general, and includes some Bogoliubov transformations as special cases.

First, let us recall that the Cuntz algebra $\mathcal{O}_d$ is a C^* -algebra generated by s_i , $i = 1, 2, \dots, d$, which satisfy the following relations:²⁾

$$s_i^* s_j = \delta_{i,j} I, \quad (2)$$

$$\sum_{i=1}^d s_i s_i^* = I. \quad (3)$$

We often use the brief description such as $s_{i_1 i_2 \dots i_m; j_n \dots j_2 j_1} \equiv s_{i_1} s_{i_2} \dots s_{i_m} s_{j_n}^* \dots s_{j_2}^* s_{j_1}^*$, $m+n \geq 1$. The $U(1)$ -invariant subalgebra $\mathcal{O}_d^{U(1)}$ of $\mathcal{O}_d$ is given by a linear space spanned by $s_{i_1 \dots i_m; j_m \dots j_1}$, $m = 1, 2, \dots$. We consider an automorphism α_u of $\mathcal{O}_d$ obtained from a natural $U(d)$ action α as follows:

$$\begin{aligned} \alpha : U(d) &\curvearrowright \mathcal{O}_d, \\ \alpha_u(s_i) &\equiv \sum_{k=1}^d s_k u_{ki}, \quad u \in U(d), \quad i = 1, 2, \dots, d. \end{aligned} \quad (4)$$

Indeed, from the equality

$$\alpha_{u_1} \circ \alpha_{u_2} = \alpha_{u_1 u_2}, \quad u_1, u_2 \in U(d), \quad (5)$$

α becomes a $U(d)$ action on $\mathcal{O}_d$. Since $\alpha_u \circ \gamma_z = \gamma_z \circ \alpha_u$, restriction of α_u to $\mathcal{O}_d^{U(1)}$ gives an automorphism of $\mathcal{O}_d^{U(1)}$:

$$\begin{aligned} \alpha_u|_{\mathcal{O}_d^{U(1)}} &\in \text{Aut } \mathcal{O}_d^{U(1)}, \\ \alpha_u(s_{i_1 \dots i_m; j_m \dots j_1}) &= \sum_{k_1, \dots, \ell_m=1}^d u_{k_1 i_1} \dots u_{k_m i_m} u_{j_1 \ell_1}^* \dots u_{j_m \ell_m}^* s_{k_1 \dots k_m; \ell_m \dots \ell_1} \in \mathcal{O}_d^{U(1)}. \end{aligned} \quad (6)$$

Using the homogeneous embedding Ψ of $\mathcal{O}_{d^r}$ ($r = 2, 3, \dots$) into $\mathcal{O}_d$ defined by³⁾

$$\begin{aligned} \Psi : \mathcal{O}_{d^r} &\hookrightarrow \mathcal{O}_d, \\ \Psi(S_i) &\equiv s_{j_1} \dots s_{j_r}, \quad i = 1, 2, \dots, d^r; \quad j_1, \dots, j_r = 1, 2, \dots, d, \\ i-1 &= \sum_{k=1}^r (j_k - 1) d^{k-1}, \end{aligned} \quad (7)$$

where $S_1, \dots, S_{d^r}$ and $s_1, \dots, s_d$ denote the generators of $\mathcal{O}_{d^p}$ and $\mathcal{O}_d$, respectively, it is straightforward to show that, for any $u \in U(d)$, there exists $v \in U(d^r)$ such that

$$(\alpha_u \circ \Psi)(X) = (\Psi \circ \alpha_v)(X), \quad X \in \mathcal{O}_{d^r}. \quad (8)$$

The generators of CAR are denoted by a_n , $n = 1, 2, \dots$, which satisfy

$$\{a_m, a_n\} = 0, \quad \{a_m, a_n^*\} = \delta_{m,n} I, \quad m, n = 1, 2, \dots \quad (9)$$

We give a systematic embedding of CAR into $\mathcal{O}_{2^p}$ in the following. The *standard recursive fermion system of order p* (RFS_p) in $\mathcal{O}_{2^p}$ is given by a set $SR_p = (\mathbf{a}_1, \dots, \mathbf{a}_p; \zeta_p, \varphi_p)$ defined by¹⁾

$$\mathbf{a}_i \equiv \sum_{k=1}^{2^{p-i}} \sum_{\ell=1}^{2^{i-1}} (-1)^{\sum_{m=1}^{i-1} \left[\frac{\ell-1}{2^{m-1}} \right]} s_{2^i(k-1)+\ell} s_{2^{i-1}(2k-1)+\ell}^*, \quad i = 1, 2, \dots, p, \quad (10)$$

$$\zeta_p(X) \equiv \sum_{i=1}^{2^p} (-1)^{\sum_{m=1}^p \left[\frac{i-1}{2^{m-1}} \right]} s_i X s_i^*, \quad X \in \mathcal{O}_{2^p}, \quad (11)$$

$$\varphi_p(X) \equiv \rho_{2^p}(X) \equiv \sum_{i=1}^{2^p} s_i X s_i^*, \quad X \in \mathcal{O}_{2^p}, \quad (12)$$

where s_i 's ($i = 1, 2, \dots, 2^p$) are generators of $\mathcal{O}_{2^p}$, $[x]$ denotes the largest integer not greater than x , and ρ_{2^p} is the canonical endomorphism of $\mathcal{O}_{2^p}$. Here, SR_p satisfies the following relations

$$\{\mathbf{a}_i, \mathbf{a}_j\} = 0, \quad \{\mathbf{a}_i, \mathbf{a}_j^*\} = \delta_{i,j} I, \quad i, j = 1, 2, \dots, p, \quad (13)$$

$$\{\mathbf{a}_i, \zeta_p(X)\} = 0, \quad \zeta_p(X)^* = \zeta(X^*), \quad i = 1, 2, \dots, p, \quad X \in \mathcal{O}_{2^p}, \quad (14)$$

$$\zeta_p(X) \zeta_p(Y) = \varphi_p(XY), \quad X, Y \in \mathcal{O}_{2^p}. \quad (15)$$

Then, the embedding Φ_{SR_p} of CAR into $\mathcal{O}_{2^p}$ associated with SR_p is defined by

$$\begin{aligned} \Phi_{SR_p} : \text{CAR} &\hookrightarrow \mathcal{O}_{2^p}, \\ \Phi_{SR_p}(a_{p(n-1)+i}) &= \zeta_p^{n-1}(\mathbf{a}_i), \quad i = 1, 2, \dots, p, \quad n = 1, 2, \dots \end{aligned} \quad (16)$$

For better understanding, we write explicitly $\mathbf{a}_i$ and ζ_p of SR_p for $p = 1, 2, 3$:

$$SR_1 \left\{ \begin{array}{l} \mathbf{a}_1 \equiv s_{1;2}, \\ \zeta_1(X) \equiv s_1 X s_1^* - s_2 X s_2^*, \quad X \in \mathcal{O}_2; \end{array} \right. \quad (17)$$

$$SR_2 \left\{ \begin{array}{l} \mathbf{a}_1 \equiv s_{1;2} + s_{3;4}, \\ \mathbf{a}_2 \equiv s_{1;3} - s_{2;4}, \\ \zeta_2(X) \equiv s_1 X s_1^* - s_2 X s_2^* - s_3 X s_3^* + s_4 X s_4^*, \quad X \in \mathcal{O}_4; \end{array} \right. \quad (18)$$

$$SR_3 \left\{ \begin{array}{l} \mathbf{a}_1 \equiv s_{1;2} + s_{3;4} + s_{5;6} + s_{7;8}, \\ \mathbf{a}_2 \equiv s_{1;3} - s_{2;4} + s_{5;7} - s_{6;8}, \\ \mathbf{a}_3 \equiv s_{1;5} - s_{2;6} - s_{3;7} + s_{4;8}, \\ \zeta_3(X) \equiv s_1 X s_1^* - s_2 X s_2^* - s_3 X s_3^* + s_4 X s_4^* \\ \quad - s_5 X s_5^* + s_6 X s_6^* + s_7 X s_7^* - s_8 X s_8^*, \quad X \in \mathcal{O}_8. \end{array} \right. \quad (19)$$

Using mathematical induction, it is shown that $\Phi_{SR_p}(\text{CAR})$ is identical with $\mathcal{O}_{2^p}^{U(1)}$, that is, any $s_{i_1 \dots i_m; j_m \dots j_1} \in \mathcal{O}_{2^p}^{U(1)}$ is expressed in terms of a monomial in $\Phi_{SR_p}(a_k)$ and $\Phi_{SR_p}(a_\ell^*)$ with $k, \ell \leq pm$.

From (6) with $d = 2^p$ and using the fact $\mathcal{O}_{2^p}^{U(1)} = \Phi_{SR_p}(\text{CAR})$, we have

$$\alpha_u|_{\Phi_{SR_p}(\text{CAR})} \in \text{Aut } \Phi_{SR_p}(\text{CAR}), \quad u \in U(2^p). \quad (20)$$

Hence, the automorphism α_u of $\mathcal{O}_{2^p}$ induces a $U(2^p)$ action χ on CAR as follows

$$\begin{aligned} \chi : U(2^p) &\curvearrowright \text{CAR}, \\ \chi_u(a_n) &\equiv (\Phi_{SR_p}^{-1} \circ \alpha_u \circ \Phi_{SR_p})(a_n), \quad u \in U(2^p), \quad n = 1, 2, \dots \end{aligned} \quad (21)$$

Here, the rhs of (21) is expressed in terms of a finite polynomial in a_k and a_ℓ^* with $k, \ell \leq pm$ for $p(m-1) + 1 \leq n \leq pm$. Therefore, χ_u gives a nonlinear transformation of CAR associated with $u \in U(2^p)$. It is obvious that the whole set of χ_u with $u \in U(2^p)$ denoted by

$$\text{Aut}_{U(2^p)}(\text{CAR}) \equiv \{\chi_u \mid u \in U(2^p)\} \quad (22)$$

constitutes a kind of nonlinear realization of $U(2^p)$ on CAR. Restricting (8) with $d = 2^p$ to $\Phi_{SR_p}(\text{CAR})$, we obtain the following inclusion relation

$$\text{Aut}_{U(2^p)}(\text{CAR}) \subset \text{Aut}_{U(2^{rp})}(\text{CAR}), \quad r = 2, 3, \dots \quad (23)$$

Hence, it may be possible to define some kind of inductive limit as $\text{Aut}_{U(2^\infty)}(\text{CAR})$, but we will not proceed in detail here.

Although the $U(2^p)$ action χ on CAR is defined individually for each $p \in \mathbf{N}$, it is possible to consider a product $\chi_{u_1} \circ \chi_{u_2}$ even for $u_1 \in U(2^{p_1})$ and $u_2 \in U(2^{p_2})$ with $p_1 \neq p_2$ according to its definition. From (23), this product is also given explicitly in $\text{Aut}_{U(2^q)}(\text{CAR})$ with q being a common multiple of p_1 and p_2 . Indeed, there exist $v_1, v_2 \in U(2^q)$ satisfying $\chi_{u_i} = \chi_{v_i}$, $i = 1, 2$, hence, we have $\chi_{u_1} \circ \chi_{u_2} = \chi_{v_1 v_2}$. In any way, we obtain an infinite-dimensional automorphism group of CAR by giving the whole set of such products as follows

$$\text{Aut}_U(\text{CAR}) \equiv \{\chi_{u_1} \circ \dots \circ \chi_{u_k} \mid u_i \in U(2^{p_i}), \quad i = 1, 2, \dots, k; \quad p_i, k = 1, 2, \dots\}. \quad (24)$$

Hereafter, we consider the nonlinear transformation of CAR defined by (21) more concretely. We rewrite (11) as $\zeta_p(X) = \sum_{i \in \mathcal{I}_+} s_i X s_i^* - \sum_{i \in \mathcal{I}_-} s_i X s_i^*$ with $\mathcal{I}_\pm$ being a certain division of indices $\{1, 2, \dots, 2^p\}$ satisfying $\sharp \mathcal{I}_\pm = 2^{p-1}$. Then, for any $u \in U(2^p) \cap U(2^{p-1}, 2^{p-1}) = U(2^{p-1}) \times U(2^{p-1})$, where each $U(2^{p-1})$ acts only on s_i with $i \in \mathcal{I}_\pm$

(or $i \in \mathcal{I}_-$), we have an equality $\alpha_u(\zeta_p(X)) = \zeta_p(\alpha_u(X))$, that is, ζ_p is *covariant* with respect to α_u for $u \in U(2^{p-1}) \times U(2^{p-1})$. On the other hand, if k is an odd positive integer, we have an identity $\zeta_p(X_1 \cdots X_k) = \zeta_p(X_1) \cdots \zeta_p(X_k)$ ($X_1, \dots, X_k \in \mathcal{O}_{2^p}$). Therefore, if $u \in U(2^{p-1}) \times U(2^{p-1})$ and $\alpha_u(\mathbf{a}_i)$ involves only odd order monomials in $\mathbf{a}_i$ ($i = 1, \dots, p$), $\{\alpha_u(\Phi_{SR_p}(a_{p(n-1)+i})) \mid i = 1, 2, \dots, p\}$ for each n is expressed in terms of polynomials in the same form as $\{\alpha_u(\mathbf{a}_i) \mid i = 1, 2, \dots, p\}$, that is, Φ_{SR_p} is *covariant* with respect to α_u .

For example, for $p \geq 2$, let u be the transposition of $s_{2^{p-1}+1}$ and s_{2^p} . Then, from an identity $\mathbf{a}_1 \mathbf{a}_2 \cdots \mathbf{a}_p = (-1)^{\lfloor \frac{p}{2} \rfloor} s_{1,2^p}$, $\alpha_u(\mathbf{a}_p)$ is expressed in terms of a polynomial involving a term $(-1)^{\lfloor \frac{p}{2} \rfloor} \mathbf{a}_1 \cdots \mathbf{a}_p$. Furthermore, if p is odd, not only ζ_p but also Φ_{SR_p} is covariant with respect to this α_u .

In the following, we present some examples explicitly.

- (1) $p = 1$, $u = u_1(\theta) \in SO(2) \subset U(2)$; $\alpha_{u_1(\theta)}(s_1) \equiv \cos \theta s_1 - \sin \theta s_2$, $\alpha_{u_1(\theta)}(s_2) \equiv \sin \theta s_1 + \cos \theta s_2$, ($0 \leq \theta < 2\pi$):

$$\begin{aligned} \chi_{u_1(\theta)}(a_1) &= \cos^2 \theta a_1 - \sin^2 \theta a_1^* + \sin \theta \cos \theta (I - 2a_1^* a_1) \\ &= \cos^2 \theta a_1 - \sin^2 \theta a_1^* + \sin \theta \cos \theta \exp(i\pi a_1^* a_1), \end{aligned} \quad (25)$$

$$\begin{aligned} \chi_{u_1(\theta)}(a_n) &= \prod_{k=1}^{n-1} \left(\cos 2\theta - \sin 2\theta \exp\left(i\pi \sum_{i=1}^{k-1} a_i^* a_i\right) (a_k^* - a_k) \right) \\ &\quad \times \left(\cos^2 \theta a_n - \sin^2 \theta a_n^* + \sin \theta \cos \theta \exp\left(i\pi \sum_{j=1}^n a_j^* a_j\right) \right), \quad n \geq 2, \end{aligned} \quad (26)$$

where use has been made of an identity $(a_k^* a_k)^2 = a_k^* a_k$, and $\exp(i\pi a_k^* a_k)$ being the Klein operator anticommuting with a_k . One should note that an identity $\exp(i2\pi a_k^* a_k) = I$ holds. From (23), the same nonlinear transformation is involved in $\text{Aut}_{U(2^p)}(\text{CAR})$ for any $p \in \mathbb{N}$. Indeed, it is straightforward to show it for $p = 2$.

- (2) $p = 2$, $u = u_2 \in \mathfrak{S}_4 \subset U(4)$; $\alpha_{u_2}(s_3) \equiv s_4$, $\alpha_{u_2}(s_4) \equiv s_3$, $\alpha_{u_2}(s_i) \equiv s_i$ for $i = 1, 2$:

$$\chi_{u_2}(a_1) = a_1 + (a_1^* - a_1) a_2^* a_2, \quad (27)$$

$$\chi_{u_2}(a_2) = -(a_1^* + a_1) a_2, \quad (28)$$

$$\chi_{u_2}(a_{2n-1}) = \exp\left(i\pi \sum_{k=1}^{n-1} a_{2k}^* a_{2k}\right) (a_{2n-1} + (a_{2n-1}^* - a_{2n-1}) a_{2n}^* a_{2n}), \quad n \geq 2, \quad (29)$$

$$\chi_{u_2}(a_{2n}) = -\exp\left(i\pi \sum_{k=1}^{n-1} a_{2k-1}^* a_{2k-1}\right) (a_{2n-1}^* + a_{2n-1}) a_{2n}, \quad n \geq 2. \quad (30)$$

It is straightforward to confirm that the equality $(\alpha_{u_2} \circ \alpha_{u_2})(a_n) = a_n$ is satisfied, which is the result of $u_2 = u_2^{-1}$.

(3) $p = 2$, $u = u_3 \in \mathfrak{S}_4 \subset U(4)$; $\alpha_{u_3}(s_2) \equiv s_3$, $\alpha_{u_3}(s_3) \equiv s_2$, $\alpha_{u_3}(s_i) \equiv s_i$ for $i = 1, 4$:

$$\chi_{u_3}(a_{2n-1}) = \exp\left(i\pi a_{2n-1}^* a_{2n-1}\right) a_{2n}, \quad (31)$$

$$\chi_{u_3}(a_{2n}) = \exp\left(i\pi a_{2n}^* a_{2n}\right) a_{2n-1}. \quad (32)$$

This transformation should be called a double Klein transformation which preserves anticommutativity between two annihilation (creation) operators.

(4) $p = 2$, $u = u_4(\theta) \in SO(2) \subset U(4)$; $\alpha_{u_4(\theta)}(s_1) \equiv \cos \theta s_1 - \sin \theta s_4$, $\alpha_{u_4(\theta)}(s_4) \equiv \sin \theta s_1 + \cos \theta s_4$, $(0 \leq \theta < 2\pi)$, $\alpha_{u_4(\theta)}(s_i) = s_i$ for $i = 2, 3$:

$$\chi_{u_4(\theta)}(a_{2n-1}) = \cos \theta a_{2n-1} + \sin \theta a_{2n}^*, \quad (33)$$

$$\chi_{u_4(\theta)}(a_{2n}) = -\sin \theta a_{2n-1}^* + \cos \theta a_{2n}. \quad (34)$$

This is nothing but an example of the conventional Bogoliubov transformation. Similar Bogoliubov transformations are obtained in $\text{Aut}_{U(4^r)}(\text{CAR})$ with $r \in \mathbf{N}$.

(5) $p = 3$, $u = u_5 \in \mathfrak{S}_8 \subset U(8)$; $\alpha_{u_5}(s_5) \equiv s_8$, $\alpha_{u_5}(s_8) \equiv s_5$, $\alpha_{u_5}(s_i) \equiv s_i$ for $i \neq 5, 8$:

$$\chi_{u_5}(a_{3n-2}) = a_{3n-2} - a_{3n-2} a_{3n}^* a_{3n} + (I - 2a_{3n-2}^* a_{3n-2}) a_{3n-1}^* a_{3n}^* a_{3n}, \quad (35)$$

$$\chi_{u_5}(a_{3n-1}) = a_{3n-1} - a_{3n-2} a_{3n}^* a_{3n} - a_{3n-2}^* (I - 2a_{3n-1}^* a_{3n-1}) a_{3n}^* a_{3n}, \quad (36)$$

$$\begin{aligned} \chi_{u_5}(a_{3n}) = & -(a_{3n-2} a_{3n-1} + a_{3n-2}^* a_{3n-2} + a_{3n-1}^* a_{3n-1} \\ & + a_{3n-2}^* a_{3n-1} - 2a_{3n-2}^* a_{3n-2} a_{3n-1}^* a_{3n-1}) a_{3n}. \end{aligned} \quad (37)$$

This is an example mentioned above for the case $p = 3$. Here, $\chi_{u_5}(a_{3n})$ involves a term $-a_{3n-2} a_{3n-1} a_{3n}$, and Φ_{SR_3} is covariant with respect to α_{u_5} .

In conclusion, we summarize the properties of the above nonlinear transformations constituting $\text{Aut}_U(\text{CAR})$:

- (i) For any $\chi \in \text{Aut}_U(\text{CAR})$, $\chi(a_n)$ is expressed in terms of a finite polynomial in generators of CAR.
- (ii) Action of any compact group on CAR is realized by certain elements of $\text{Aut}_U(\text{CAR})$.
- (iii) Through a nonlinear transformation of CAR, particle numbers are not preserved in general.
- (iv) A subset of $\text{Aut}_U(\text{CAR})$ consisting of all transformations, which keep the Fock vacuum invariant, constitutes a nontrivial subgroup of $\text{Aut}_U(\text{CAR})$.
- (v) In general, $\chi \in \text{Aut}_U(\text{CAR})$ is an outer automorphism, that is, it is *not* expressed by the adjoint operation of a unitary element in CAR as seen by checking the action of χ on the Fock space. Restriction of the permutation representation of Cuntz algebra $\mathcal{O}_{2^p}$ to $\Phi_{SR_p}(\text{CAR})$ determines the change of representations of CAR through the nonlinear transformation.

References

- 1) M. Abe and K. Kawamura, "*Recursive Fermion System in Cuntz Algebra. I — Embeddings of Fermion Algebra into Cuntz Algebra —*", Preprint RIMS-1332, math-ph/0110003.
- 2) J. Cuntz, Comm. Math. Phys. **57**, 173, (1977).
- 3) M. Abe and K. Kawamura, "*Recursive Fermion System in Cuntz Algebra. II — Endomorphism, Automorphism and Branching of Representation —*", in preparation.